

Express Mail No. EV 272754553 US

**PREPARATION OF SUBSTITUTED ISOSERINE ESTERS
USING β -LACTAMS AND METAL OR AMMONIUM ALKOXIDES**


REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation of U.S. Serial No.
10/194,343, filed July 12, 2002, which is a continuation of U.S.
Serial No. 09/516,870, filed on March 2, 2000, which is a
continuation of U.S. Serial No. 08/938,075, filed on September
26, 1997, now U.S. Patent No. 6,124,481, which is a division of
U.S. Serial No. 08/476,357, filed on June 7, 1995, now U.S.
10 Patent 5,717,115, which is a division of U.S. Serial No.
08/034,247, filed on March 22, 1993, now U.S. Patent No.
5,430,160.

BACKGROUND OF THE INVENTION

15 Three esters of N-acyl phenyl isoserine, taxol, taxotere and
cephalomannine have been found to possess significant properties
as antitumor agents. This application describes a process for
the preparation of N-acyl, N-sulfonyl and N-phosphoryl
substituted isoserine esters, in general and to a semi-synthesis
for the preparation of taxane derivatives such as taxol, taxotere
20 and other biologically active derivatives involving the use of
metal alkoxides and β -lactams, in particular.

The taxane family of terpenes, of which taxol is a member,
has attracted considerable interest in both the biological and
chemical arts. Taxol is a promising cancer chemotherapeutic
25 agent with a broad spectrum of antileukemic and tumor-inhibiting
activity. Taxol has the following structure:




wherein Ac is acetyl. Because of this promising activity, taxol is currently undergoing clinical trials in both France and the United States.

5 The supply of taxol for these clinical trials is presently being provided by the bark from Taxus brevifolia (Western Yew). However, taxol is found only in minute quantities in the bark of these slow growing evergreens, causing considerable concern that the limited supply of taxol
10 will not meet the demand. Consequently, chemists in recent years have expended their energies in trying to find a viable synthetic route for the preparation of taxol. So far, the results have not been entirely satisfactory.

15 One synthetic route that has been proposed is directed to the synthesis of the tetracyclic taxane nucleus from commodity chemicals. A synthesis of the taxol congener taxusin has been reported by Holton, et al. in JACS 110, 6558 (1988). Despite the progress made in this approach, the final total synthesis of taxol is, nevertheless, likely to be
20 a multi-step, tedious, and costly process.


25 A semi-synthetic approach to the preparation of taxol has been described by Greene, et al. in JACS 110, 5917 (1988), and involves the use of a congener of taxol, 10-deacetyl baccatin III which has the structure of formula II shown below:




10-deacetyl baccatin III is more readily available than taxol since it can be obtained from the needles of Taxus baccata.

According to the method of Greene et al., 10-deacetyl
5 baccatin III is converted to taxol by attachment of the C-10 acetyl group and by attachment of the C-13 β -amido ester side chain through the esterification of the C-13 alcohol with a β -amido carboxylic acid unit. Although this approach
10 requires relatively few steps, the synthesis of the β -amido carboxylic acid unit is a multi-step process which proceeds in low yield, and the coupling reaction is tedious and also proceeds in low yield. However, this coupling reaction is a
15 key step which is required in every contemplated synthesis of taxol or biologically active derivative of taxol, since it has been shown by Wani, et al. in JACS 93, 2325 (1971) that the presence of the β -amido ester side chain at C13 is required for anti-tumor activity.

More recently, it has been reported in Colin et al. U.S. Patent No. 4,814,470 that taxanes corresponding to the
20 following formula III, have an activity significantly greater than that of taxol (I).




According to Colin et al., U.S. Patent 4,418,470, the products of general formula (III) are obtained by the action of the sodium salt of tert-butyl N-chlorocarbamate on a product of general formula:




preparation of the product of general formula (IV) can be converted to a product of general formula (III).

In an effort to improve upon the Colin et al. process, Denis et al. disclose a different process for preparing derivatives of baccatin III or of 10-deacetyl-baccatin III of general formula



in which R' denotes hydrogen or acetyl wherein an acid of general formula:



in which R₁ is a hydroxy-protecting group, is condensed with a taxane derivative of general formula:



5 However, this method employs relatively harsh conditions,
proceeds with poor conversion, and provides less than optimal
yields.

10

15

20


25

permitting the use of a racemic mixture of side chain precursor, eliminating the need for the expensive, time-consuming process of separating the precursor into its respective enantiomeric forms. The reaction additionally proceeds at a faster rate than previous processes, thus permitting the use of less side-chain precursor than has been required by such previous processes.


SUMMARY OF THE INVENTION

Among the objects of the present invention, therefore, is the provision of a process for the preparation of N-acyl, N-sulfonyl and N-phosphoryl esters of isoserine; the provision of a side chain precursor for the synthesis of taxane derivatives; the provision of a process for the attachment of the side chain precursor in relatively high yield to provide an intermediate which is readily converted to the desired taxane derivative; and the provision of such a process which is highly diastereoselective.


In accordance with the present invention, a process is provided for the preparation of isoserine esters having the formula



comprising reacting a β -lactam with an alkoxide, the β -lactam having the formula



and the alkoxide having the formula



wherein

R_1 is $-\text{OR}_6$, $-\text{SR}_7$, or $-\text{NR}_8\text{R}_9$;

R_2 is hydrogen, alkyl, alkenyl, alkynyl, aryl, or heteroaryl;

R_3 and R_4 are independently hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, acyl or heterosubstituted alkyl, alkenyl, alkynyl, aryl or heteroaryl, provided,

however, that R_3 and R_4 are not both acyl;

R_5 is $-\text{COR}_{10}$, $-\text{COOR}_{10}$, $-\text{COSR}_{10}$, $-\text{CONR}_8\text{R}_{10}$, or $-\text{SO}_2\text{R}_{11}$,

R_6 is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, hydroxy protecting group, or a functional group which increases the water solubility of the taxane derivative,

R_7 is alkyl, alkenyl, alkynyl, aryl, heteroaryl, or sulfhydryl protecting group,

R_8 is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, or heterosubstituted alkyl, alkenyl, alkynyl, aryl or heteroaryl;

R_9 is an amino protecting group;

R_{10} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, or heterosubstituted alkyl, alkenyl alkynyl, aryl or heteroaryl;


R_{11} is alkyl, alkenyl, alkynyl, aryl, heteroaryl, $-\text{OR}_{10}$, or $-\text{NR}_8\text{R}_{14}$;

R_{14} is hydrogen, alkyl, alkenyl, alkynyl, aryl, or heteroaryl;

E_1 , E_2 and E_3 are independently hydrogen, hydrocarbon or cyclic, provided, at least one of E_1 , E_2 and E_3 is other than hydrogen; and

M comprises ammonium or is a metal.

In accordance with another aspect of the present invention, the alkoxide and β -lactam are selected so as to provide a process for preparing taxol, taxotere and other
 5 biologically active taxane derivatives having the following structural formula:



wherein

- $R_1 - R_{14}$ are as previously defined,
- 10 R_{15} is hydrogen or together with R_{16} forms an oxo,
- R_{16} is hydrogen, $-OCOR_{29}$, hydroxy, or protected hydroxy, or together with R_{15} forms an oxo;
- R_{17} is hydrogen or together with R_{18} forms an oxo,
- R_{18} is hydrogen, hydroxy, protected hydroxy,
- 15 acyloxy, or together with R_{17} forms an oxo;
- R_{19} is hydrogen or together with R_{20} forms an oxo,
- R_{20} is hydrogen, halogen, protected hydroxy, $-OR_{28}$, or together with R_{19} forms an oxo;
- R_{21} is hydrogen or together with R_{22} forms an oxo,
- 20 R_{22} is hydrogen, hydroxy, protected hydroxy, acyloxy, together with R_{21} forms an oxo, or together with R_{23} and the carbon atoms to which they are attached form an oxetane ring;
- R_{23} is hydrogen, together with R_{24} forms an oxo, or
- 25 together with R_{22} and the carbon atoms to which they are

attached form an oxetane ring;

R_{24} is hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, cyano, hydroxy, $-\text{OCOR}_{30}$, or together with R_{23} forms an oxo, oxirane or methylene;

5 R_{25} is hydrogen, hydroxy, or $-\text{OCOR}_{31}$;


R_{26} is hydrogen or taken together with R_{25} forms an oxo;

R_{27} is hydrogen, hydroxy, protected hydroxy;


10 R_{28} is hydrogen, acyl, hydroxy protecting group or a functional group which increases the solubility of the taxane derivative; and

R_{29} , R_{30} , and R_{31} are independently hydrogen, alkyl, alkenyl, alkynyl, monocyclic aryl or monocyclic heteroaryl.

15 Briefly, therefore, the taxane derivatives are prepared by reacting a β -lactam (2) with an alkoxide having the bi-, tri- or tetracyclic taxane nucleus to form a β -amido ester intermediate. The intermediate is then converted to the taxane derivative. β -lactam (2) has the general formula:



20 wherein $R_1 - R_5$ are as previously defined. The alkoxide preferably has the tricyclic taxane nucleus corresponding to the general formula:



(3)

wherein M and R₁₅-R₂₇ are as previously defined. Most preferably, the alkoxide has the tetracyclic taxane nucleus corresponding to alkoxide (3) wherein R₂₂ and R₂₃ together form an oxetane ring.

Other objects and features of this invention will be in part apparent and in part pointed out hereinafter.

DETAILED DESCRIPTION

As used herein "Ar" means aryl; "Ph" means phenyl; "Ac" means acetyl; "Et" means ethyl; "R" means alkyl unless otherwise defined; "Bu" means butyl; "Pr" means propyl; "TES" means triethylsilyl; "TMS" means trimethylsilyl; "TPAP" means tetrapropylammonium perruthenate; "DMAP" means p-dimethylamino pyridine; "DMF" means dimethylformamide; "LDA" means lithium diisopropylamide; "LAH" means lithium aluminum hydride; "Red-Al" means sodium bis(2-methoxyethoxy) aluminum hydride; "AIBN" means azo-(bis)-isobutyronitrile "10-DAB" means 10-desacetylbaicatin III; protected hydroxy means -OR wherein R is a hydroxy protecting group; sulfhydryl protecting group" includes, but is not limited to, hemithioacetals such as 1-ethoxyethyl and methoxymethyl, thioesters, or thiocarbonates; "amine protecting group" includes, but is not limited to, carbamates, for example, 2,2,2-trichloroethylcarbamate or tertbutylcarbamate; and "hydroxy protecting group" includes, but is not limited to,

ethers such as methyl, t-butyl, benzyl, p-methoxybenzyl, p-nitrobenzyl, allyl, trityl, methoxymethyl, methoxyethoxymethyl, ethoxyethyl, tetrahydropyranyl, tetrahydrothiopyranyl, and trialkylsilyl ethers such as trimethylsilyl ether, triethylsilyl ether, dimethylarylsilyl ether, triisopropylsilyl ether and t-butyl dimethylsilyl ether; esters such as benzoyl, acetyl, phenylacetyl, formyl, mono-, di-, and trihaloacetyl such as chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl; and carbonates including but not limited to alkyl carbonates having from one to six carbon atoms such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl; isobutyl, and n-pentyl; alkyl carbonates having from one to six carbon atoms and substituted with one or more halogen atoms such as 2,2,2-trichloroethoxymethyl and 2,2,2-trichloroethyl; alkenyl carbonates having from two to six carbon atoms such as vinyl and allyl; cycloalkyl carbonates have from three to six carbon atoms such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl; and phenyl or benzyl carbonates optionally substituted on the ring with one or more C_{1-5} alkoxy, or nitro. Other hydroxyl, sulfhydryl and amine protecting groups may be found in "Protective Groups in Organic Synthesis" by T. W. Greene, John Wiley and Sons, 1981.

The alkyl groups described herein, either alone or with the various substituents defined hereinabove are preferably lower alkyl containing from one to six carbon atoms in the principal chain and up to 15 carbon atoms. They may be straight or branched chain and include methyl, ethyl, propyl, isopropyl, butyl, hexyl and the like.

The alkenyl groups described herein, either alone or with the various substituents defined hereinabove are preferably lower alkenyl containing from two to six carbon

atoms in the principal chain and up to 15 carbon atoms. They may be straight or branched chain and include ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, hexenyl, and the like.

5 The alkynyl groups described herein, either alone or with the various substituents defined hereinabove are preferably lower alkynyl containing from two to six carbon atoms in the principal chain and up to 15 carbon atoms. They may be straight or branched chain and include ethynyl,
10 propynyl, butynyl, isobutynyl, hexynyl, and the like.

 The aryl moieties described herein, either alone or with various substituents, contain from 6 to 15 carbon atoms and include phenyl. Substituents include alkanoxy, protected hydroxy, halogen, alkyl, aryl, alkenyl, acyl, acyloxy, nitro,
15 amino, amido, etc. Phenyl is the more preferred aryl.


 The heteroaryl moieties described herein, either alone or with various substituents, contain from 5 to 15 atoms and include, furyl, thienyl, pyridyl and the like. Substituents include alkanoxy, protected hydroxy, halogen,
20 alkyl, aryl, alkenyl, acyl, acyloxy, nitro, amino, and amido.

 The acyloxy groups described herein contain alkyl, alkenyl, alkynyl, aryl or heteroaryl groups.

 The heterosubstituents of the heterosubstituted alkyl, alkenyl, alkynyl, aryl, and heteroaryl moieties
25 described herein, contain nitrogen, oxygen, sulfur, halogens and/or one to six carbons, and include lower alkoxy such as methoxy, ethoxy, butoxy, halogen such as chloro or fluoro, and nitro.


 The present invention is directed to a process for
30 preparing substituted isoserine esters, in general, and taxol, taxotere and other taxane derivatives which are biologically active using β -lactam (2), the structure of

which is depicted hereinbelow:




wherein R_1 , R_2 , R_3 , R_4 and R_5 are as previously defined.

- In accordance with the present invention, R_5 of β -lactam (2) is preferably $-\text{COR}_{10}$ or $-\text{COOR}_{10}$ with R_{10} being lower alkyl, aryl, heteroaryl (such as furyl or thienyl), or substituted phenyl, and most preferably phenyl, methyl, ethyl, tert-butyl, or



- 10 wherein X is Cl, Br, F, $\text{CH}_3\text{O}-$, or NO_2- . Preferably R_2 and R_4 are hydrogen or lower alkyl. R_3 is preferably aryl, most preferably, naphthyl, phenyl,



- 15 wherein X is as previously defined, Me is methyl and Ph is phenyl. Preferably, R_1 is selected from $-\text{OR}_6$, $-\text{SR}_7$ or $-\text{NR}_8\text{R}_9$ wherein R_6 , R_7 and R_9 , are hydroxy, sulfhydryl, and amine


protecting groups, respectively, and R_8 is hydrogen, alkyl, alkenyl, alkynyl, aryl, or heteroaryl. Most preferably, R_1 is $-OR_6$ wherein R_6 is triethylsilyl ("TES"), 1-ethoxyethyl ("EE") or 2,2,2-trichloroethoxymethyl.

5 As noted above, R_1 of β -lactam (2) may be $-OR_6$ with R_6 being alkyl, acyl, ethoxyethyl ("EE"), triethylsilyl ("TES"), 2,2,2-trichloroethoxymethyl, or other hydroxyl protecting group such as acetals and ethers, i.e., methoxymethyl ("MOM"), benzyloxymethyl; esters, such as
10 acetates; carbonates, such as methyl carbonates; and alkyl and aryl silyl such as triethylsilyl, trimethylsilyl, dimethyl-t-butylsilyl, dimethylarylsilyl, dimethyl-heteroarylsilyl, and triisopropylsilyl, and the like. A variety of protecting groups for the hydroxyl group and the
15 synthesis thereof may be found in "Protective Groups in Organic Synthesis" by T. W. Greene, John Wiley and Sons, 1981. The hydroxyl protecting group selected should be easily removed under conditions that are sufficiently mild, e.g., in 48% HF, acetonitrile, pyridine, or 0.5% HCl/water/
20 ethanol, and/or zinc, acetic acid so as not to disturb the ester linkage or other substituents of the taxol intermediate. However, R_6 is preferably triethylsilyl, 1-ethoxyethyl or 2,2,2-trichloroethoxymethyl, and most preferably triethylsilyl.


25 Since β -lactam (2) has several asymmetric carbons, it is known to those skilled in the art that the compounds of the present invention having asymmetric carbon atoms may exist in diastereomeric, racemic, or optically active forms. All of these forms are contemplated within the scope of this
30 invention. More specifically, the present invention includes enantiomers, diastereomers, racemic mixtures, and other mixtures thereof.

β -lactam (2) can be prepared from readily available materials, as is illustrated in schemes A and B below:

Scheme A



Scheme B



reagents: (a) triethylamine, CH_2Cl_2 , 25°C , 18h; (b) 4 equiv ceric ammonium nitrate, CH_3CN , -10°C , 10 min; (c) KOH, THF,

H₂O, 0°C, 30 min; (d) ethyl vinyl ether, THF, toluene sulfonic acid (cat.), 0°C, 1.5h; (e) n-butyllithium, ether, -78°C, 10 min; benzoyl chloride, -78°C, 1h; (f) lithium diisopropyl amide, THF -78°C to -50°C; (g) lithium
5 hexamethyldisilazide, THF -78°C to 0°C; (h) THF, -78°C to 25°C, 12h.


The starting materials are readily available. In scheme A, α -acetoxy acetyl chloride is prepared from glycolic acid, and, in the presence of a tertiary amine, it
10 cyclocondenses with imines prepared from aldehydes and p-methoxyaniline to give 1-p-methoxyphenyl-3-acyloxy-4-arylazetidin-2-ones. The p-methoxyphenyl group can be readily removed through oxidation with ceric ammonium nitrate, and the acyloxy group can be hydrolyzed under
15 standard conditions familiar to those experienced in the art to provide 3-hydroxy-4-arylazetidin-2-ones. The 3-hydroxyl group is protected with 1-ethoxyethyl, but may be protected with variety of standard protecting groups such as the triethylsilyl group or other trialkyl (or aryl) silyl groups.
20 In Scheme B, ethyl- α -triethylsilyloxyacetate is readily prepared from glycolic acid.

The racemic β -lactams may be resolved into the pure enantiomers prior to protection by recrystallization of the corresponding 2-methoxy-2-(trifluoromethyl) phenylacetic
25 esters. However, the reaction described hereinbelow in which the β -amido ester side chain is attached has the advantage of being highly diastereoselective, thus permitting the use of a racemic mixture of side chain precursor.

The 3-(1-ethoxyethoxy)-4-phenylazetidin-2-one of
30 scheme A and the 3-(1-triethylsilyloxy)-4-phenylazetidin-2-one of scheme B can be converted to β -lactam (2), by treatment with a base, preferably n-butyllithium, and an acyl


chloride, alkylchloroformate, sulfonyl chloride, phosphinyl chloride or phosphoryl chloride at -78 °C or below.

The process of the present invention is particularly useful for the esterification of mono- or polycyclic alkoxides represented by the formula



in which E_1 , E_2 and the carbon to which they are attached define a carbocyclic and/or heterocyclic skeleton which may be mono- or polycyclic and E_3 is hydrogen or hydrocarbon, preferably lower alkyl. Most preferably, the carbocyclic and/or heterocyclic skeleton comprises about 6 to 20 atoms and the hetero atoms are oxygen. The cyclic skeleton may be hydrocarbon and/or heterosubstituted with heterosubstituents including, for example, esters, ethers, amines, alcohols, protected alcohols, carbonyl groups, halogens, oxygen, substituted oxygen or substituted nitrogen.


When the alkoxides have the bi-, tri- or tetracyclic taxane nucleus, the process of the present invention may advantageously be used to prepare taxane derivatives, many of which have been found to have significant biological activity. As used herein, an alkoxide having the bicyclic taxane nucleus has the carbocyclic skeleton corresponding to rings A and B of metal alkoxide (3):



wherein M and R_{15} - R_{27} are as previously defined. An alkoxide having the tricyclic taxane nucleus has the carbocyclic skeleton corresponding to rings A, B and C of metal alkoxide (3). An alkoxide having the tetracyclic taxane nucleus has carbocyclic rings A, B and C of metal alkoxide (3) and the oxetane ring defined by R_{22} , R_{23} , and the carbons to which they are attached.


Substituent, M, of alkoxide 3 is a metal or comprises ammonium. The metal may be a Group IA, IIA, transition (including lanthanides and actinides), IIB, IIIA, IVA, VA, or VIA metal (CAS version). The ammonium comprising substituent is preferably tetraalkylammonium and the alkyl component of the tetraalkylammonium substituent is preferably C_1 - C_{10} alkyl such as methyl or butyl.

Alkoxides (3) are prepared by reacting an alcohol having two to four rings of the taxane nucleus and a C-13 hydroxyl group with an organometallic compound, zinc chloride dimethoxyethane complex, anhydrous cadmium chloride, or a tetraalkylammonium halide such as tetrabutylammonium chloride in a suitable solvent. Preferably, the alcohol is a derivative of baccatin III or 10-deacetyl baccatin III having the structure:



wherein R_{16} - R_{20} , R_{24} , and R_{26} are as previously defined. More preferably, R_{16} is protected hydroxy or $-OCOR_{29}$, R_{19} is hydrogen, R_{20} is hydrogen, halogen or protected hydroxy, R_{24} is acetoxy, and R_{26} is benzoyloxy. Most preferably, the alcohol is a protected baccatin III, in particular, 7-O-triethylsilyl baccatin III (which can be obtained as described by Greene, et al. in JACS 110, 5917 (1988) or by other routes) or 7,10-bis-O-triethylsilyl baccatin III.




As reported in Greene et al., 10-deacetyl baccatin III is converted to 7-O-triethylsilyl-10-deacetyl baccatin III according to the following reaction scheme:



Under what is reported to be carefully optimized conditions, 10-deacetyl baccatin III is reacted with 20 equivalents of $(C_2H_5)_3SiCl$ at $23^\circ C$ under an argon atmosphere for 20 hours in the presence of 50 ml of pyridine/mmol of 10-deacetyl baccatin III to provide 7-triethylsilyl-10-deacetyl baccatin

III (6a) as a reaction product in 84-86% yield after purification. The reaction product is then acetylated with 5 equivalents of CH_3COCl and 25 mL of pyridine/mmol of (6a) at 0 °C under an argon atmosphere for 48 hours to provide 86% yield of 7-O-triethylsilyl baccatin III (6b). Greene, et al. in JACS 110, 5917 at 5918 (1988).



Alternatively, 7-triethylsilyl-10-deacetyl baccatin III (6a) can be protected at C-10 oxygen with an acid labile hydroxyl protecting group. For example, treatment of (6a) with n-butyllithium in THF followed by triethylsilyl chloride (1.1 mol equiv.) at 0°C gives 7,10-bis-O-triethylsilyl baccatin III (6c) in 95% yield. Also, (6a) can be converted to 7-O-triethylsilyl-10-(1-ethoxyethyl) baccatin III (6d) in 90% yield by treatment with excess ethyl vinyl ether and a catalytic amount of methane sulfonic acid. These preparations are illustrated in the reaction scheme below.



5 7-O-triethylsilyl baccatin III (6b), 7,10-bis-O-triethylsilyl baccatin III (6c), or 7-O-triethylsilyl-10-(1-ethoxyethyl) baccatin III (6d) is reacted with an organometallic compound such as n-butyllithium in a solvent such as tetrahydrofuran (THF), to form the metal alkoxide

10 13-O-lithium-7-O-triethylsilyl baccatin III (7b)


13-O-lithium-7,10-bis-O-triethylsilyl baccatin III (7c), or 13-O-lithium-7-O-triethylsilyl-10-(1-ethoxyethyl) baccatin III (7d) as shown in the following reaction scheme:



- (7b) Z = $-\text{OCOCH}_3$
 (7c) Z = $-\text{OSi}(\text{C}_2\text{H}_5)_3$
 (7d) Z = $-\text{OEE}$

As illustrated in the following reaction scheme, a suitable metal alkoxide of the present invention such as 13-O-lithium-7-O-triethylsilyl baccatin III derivative (7b, 7c, or 7d) reacts with a β -lactam of the present invention to provide an intermediate (8b, 8c, or 8d) in which the C-7

hydroxyl group is protected with a triethylsilyl or 1-ethoxyethyl group.




Intermediate compound (8b) readily converts to taxol when R₁ is -OR₆, R₂ and R₃ are hydrogen, R₄ is phenyl, R₅ is benzoyl and R₆ is a hydroxy protecting group such as triethylsilyl. Intermediate compound (8c) readily converts to taxotere when R₁ is -OR₆, R₂ and R₃ are hydrogen, R₄ is phenyl, R₅ is tertbutoxycarbonyl and R₆ is a hydroxy protecting group such as triethylsilyl. Intermediate compound (8d) readily converts to 10-deacetyl taxol when R₁ is -OR₆, R₂ and R₃ are hydrogen, R₄ is phenyl, R₅ is benzoyl, and R₆ is a hydroxy protecting group such as triethylsilyl. Intermediate compounds (8b, 8c and 8d) may be converted to the indicated compounds by hydrolyzing the triethylsilyl and 1-ethoxyethyl groups under mild conditions so as not to disturb the ester linkage or the taxane derivative substituents.

- 8b $\xrightarrow{\text{HF, C}_5\text{H}_5\text{N, CH}_3\text{CN}}$ TAXOL
- 5 8c $\xrightarrow{\text{HF, C}_5\text{H}_5\text{N, CH}_3\text{CN}}$ TAXOTERE
- 10 8d $\xrightarrow{0.1\% \text{ HCl, EtOH}}$ 10-DEACETYL TAXOL

Both the conversion of the alcohol to the metal alkoxide and the ultimate synthesis of the taxol can take place in the same reaction vessel. Preferably, the β -lactam is added to the reaction vessel after formation therein of the metal alkoxide.

The organometallic compound n-butyllithium is preferably used to convert the alcohol to the corresponding metal alkoxide, but other sources of metallic substituent such as lithium diisopropyl amide, other lithium or magnesium amides, ethylmagnesium bromide, methylmagnesium bromide, other organolithium compounds, other organomagnesium compounds, organosodium, organotitanium, organozirconium, organozinc, organocadmium or organopotassium or the corresponding amides may also be used. Organometallic compounds are readily available, or may be prepared by available methods including reduction of organic halides with metal. Lower alkyl halides are preferred. For example, butyl bromide can be reacted with lithium metal in diethyl ether to give a solution of n-butyllithium in the following manner:



Alternatively, the lithium alkoxide may be induced to undergo exchange with metal halides to form alkoxides of aluminum, boron, cerium, calcium, zirconium or zinc.

Although THF is the preferred solvent for the
5 reaction mixture, other ethereal solvents, such as dimethoxyethane, or aromatic solvents may also be suitable. Certain solvents, including some halogenated solvents and some straight-chain hydrocarbons in which the reactants are too poorly soluble, are not suitable. Other solvents are not
10 appropriate for other reasons. For example, esters are not appropriate for use with certain organometallic compounds such as n-butyllithium due to incompatibility therewith.

Although the reaction scheme disclosed herein is directed to the synthesis of certain taxol derivatives, it
15 can be used with modifications in either the β -lactam or the tetracyclic metal alkoxide. Therefore, alkoxides other than 13-O-lithium-7-O-triethylsilyl baccatin III may be used to form a taxol or other taxanes according to the method of this invention. The β -lactam and the tetracyclic alkoxide can be
20 derived from natural or unnatural sources, to prepare other synthetic taxols, taxol derivatives, 10-deacetyltaxols, and the enantiomers and diastereomers thereof contemplated within the present invention.

The process of the invention also has the important
25 advantage of being highly diastereoselective. Therefore racemic mixtures of the side chain precursors may be used. Substantial cost savings may be realized because there is no need to resolve racemic β -lactams into their pure enantiomers. Additional cost savings may be realized because
30 less side chain precursor, e.g., 60-70% less, is required relative to prior processes.

The water solubility of compounds of formula (3) may be improved by modification of the C2' and/or C7 substituents. For instance, water solubility may be increased if R_1 is $-OR_6$ and R_{20} is $-OR_{28}$, and R_6 and R_{28} are independently hydrogen or $-COGCOR^1$ wherein

G is ethylene, propylene, $-\text{CH}=\text{CH}-$, 1,2-cyclohexane, or 1,2-phenylene,

$R^1 = \text{OH base}, \text{NR}^2\text{R}^3, \text{OR}^3, \text{SR}^3, \text{OCH}_2\text{CONR}^4\text{R}^5, \text{OH}$

$R^2 = \text{hydrogen, methyl}$

$R^3 = (\text{CH}_2)_n\text{NR}^6\text{R}^7; (\text{CH}_2)_n\text{N}^+\text{R}^6\text{R}^7\text{R}^8\text{X}^0$


$n = 1 \text{ to } 3$

$R^4 = \text{hydrogen, lower alkyl containing 1 to 4 carbons}$

$R^5 = \text{hydrogen, lower alkyl containing 1 to 4 carbons, benzyl, hydroxyethyl, } \text{CH}_2\text{CO}_2\text{H, dimethylaminoethyl}$

$R^6\text{R}^7 = \text{lower alkyl containing 1 or 2 carbons, benzyl or } R^6 \text{ and}$

$R^7 \text{ together with the nitrogen atom of } \text{NR}^6\text{R}^7 \text{ form the following rings}$



$R^8 = \text{lower alkyl containing 1 or 2 carbons, benzyl}$

$\text{X}^0 = \text{halide}$

base = $\text{NH}_3, (\text{HOC}_2\text{H}_4)_3\text{N}, \text{N}(\text{CH}_3)_3, \text{CH}_3\text{N}(\text{C}_2\text{H}_4\text{OH})_2,$

$\text{NH}_2(\text{CH}_2)_6\text{NH}_2, \text{N-methylglucamine, NaOH, KOH.}$


The preparation of compounds in which X_1 or X_2 is $-\text{COGCOR}^1$ is set forth in Hangwitz U.S. Patent 4,942,184 which is incorporated herein by reference.

Alternatively, water solubility may be increased when R_1 is $-\text{OR}_6$ and R_6 is a radical having the formula $-\text{COCX}=\text{CHX}$ or $-\text{COX}-\text{CHX}-\text{CHX}-\text{SO}_2\text{O}-\text{M}$ wherein X is hydrogen, alkyl or aryl and M is hydrogen, alkaline metal or an ammonio group as described in Kingston et al., U.S. Patent No. 5,059,699 (incorporated herein by reference).


Taxanes having alternative C9 keto substituent may be prepared by selectively reduction to yield the corresponding C9 β -hydroxy derivative. The reducing agent is preferably a borohydride and, most preferably, tetrabutylammoniumborohydride (Bu_4NBH_4) or triacetoxyborohydride.

As illustrated in Reaction Scheme 1, the reaction of baccatin III with Bu_4NBH_4 in methylene chloride yields 9-desoxo-9 β -hydroxybaccatin III 5. After the C7 hydroxy group is protected with the triethylsilyl protecting group, for example, a suitable side chain may be attached to 7-protected-9 β -hydroxy derivative 6 as elsewhere described herein. Removal of the remaining protecting groups thus yields 9 β -hydroxy-desoxo taxol or other 9 β -hydroxytetracyclic taxane having a C13 side chain.


REACTION SCHEME 1




Alternatively, the C13 hydroxy group of 7-protected-9 β -hydroxy derivative 6 may be protected with trimethylsilyl or other protecting group which can be selectively removed relative to the C7 hydroxy protecting group as illustrated in Reaction Scheme 2, to enable further selective manipulation of the various substituents of the taxane. For example, reaction of 7,13-protected-9 β -hydroxy derivative 7 with KH causes the acetate group to migrate from C10 to C9 and the hydroxy group to migrate from C9 to C10, thereby yielding 10-desacetyl derivative 8. Protection of the C10 hydroxy group of 10-desacetyl derivative 8 with triethylsilyl yields derivative 9. Selective removal of the C13 hydroxy protecting group from derivative 9 yields derivative 10 to which a suitable side chain may be attached as described above.

REACTION SCHEME 2


As shown in Reaction Scheme 3, 10-oxo derivative 11 can be provided by oxidation of 10-desacetyl derivative 8. Thereafter, the C13 hydroxy protecting group can be selectively removed followed by attachment of a side chain as described above to yield 9-acetoxy-10-oxo-taxol or other 9-acetoxy-10-oxotetracyclic taxanes having a C13 side chain. Alternatively, the C9 acetate group can be selectively removed by reduction of 10-oxo derivative 11 with a reducing agent such as samarium diiodide to yield 9-desoxo-10-oxo derivative 12 from which the C13 hydroxy protecting group can be selectively removed followed by attachment of a side chain as described above to yield 9-desoxo-10-oxo-taxol or other 9-desoxo-10-oxotetracyclic taxanes having a C13 side chain.

REACTION SCHEME 3

Reaction Scheme 4 illustrates a series of reactions in which 10-DAB is used as the starting material. Reduction of 10-DAB yields pentaol 13, the C7 and C10 hydroxyl groups of which can be selectively protected with the triethylsilyl or another protecting group to produce triol 14. A C13 side chain can be attached to triol 14 as described above or, alternatively, after further modification of the tetracyclic substituents.

REACTION SCHEME 4

Taxanes having C9 and/or C10 acyloxy substituents other than acetate can be prepared using 10-DAB as a starting material as illustrated in Reaction Scheme 5. Reaction of 10-DAB with triethylsilyl chloride in pyridine yields 7-protected 10-DAB 15. The C10 hydroxy substituent of 7-protected 10-DAB 15 may then be readily acylated with any standard acylating agent to yield derivative 16 having a new C10 acyloxy substituent. Selective reduction of the C9 keto substituent of derivative 16 yields 9 β -hydroxy derivative 17 to which a C13 side chain may be attached. Alternatively, the C10 and C9 groups can be caused to migrate as set forth in Reaction Scheme 2, above.


REACTION SCHEME 5

Taxanes having alternative C2 and/or C4 esters can be prepared using baccatin III and 10-DAB as starting materials. The C2 and/or C4 esters of baccatin III and 10-DAB can be selectively reduced to the corresponding alcohol(s) using reducing agents such as LAH or Red-Al, and new esters can thereafter be substituted using standard acylating agents such as anhydrides and acid chlorides in combination with an amine such as pyridine, triethylamine, DMAP, or diisopropyl ethyl amine. Alternatively, the C2 and/or C4 alcohols may be converted to new C2 and/or C4 esters through formation of the corresponding alkoxide by


treatment of the alcohol with a suitable base such as LDA followed by an acylating agent such as an acid chloride.

Baccatin III and 10-DAB analogs having different substituents at C2 and/or C4 can be prepared as set forth in Reaction Schemes 6-10. To simplify the description, 10-DAB is used as the starting material. It should be understood, however, that baccatin III derivatives or analogs may be produced using the same series of reactions (except for the protection of the C10 hydroxy group) by simply replacing 10-DAB with baccatin III as the starting material. 9-desoxo derivatives of the baccatin III and 10-DAB analogs having different substituents at C2 and/or C4 can then be prepared by reducing the C9 keto substituent of these analogs and carrying out the other reactions described above.


In Reaction Scheme 6, protected 10-DAB **3** is converted to the triol **18** with lithium aluminum hydride. Triol **18** is then converted to the corresponding C4 ester using Cl_2CO in pyridine followed by a nucleophilic agent (e.g., Grignard reagents or alkylolithium reagents).


Scheme 6

Alternatively, deprotonation of triol **18** with LDA
5 followed by introduction of an acid chloride selectively
gives the C4 ester. For example, when acetyl chloride was
used, triol **18** was converted to 1,2 diol **4** as set forth in
Reaction Scheme 7.

Scheme 7

As set forth in Reaction Scheme 9, other C4 substituents can be provided by reacting carbonate 19 with an acid chloride and a tertiary amine to yield carbonate 22 which is then reacted with alkylolithiums or Grignard reagents to provide 10-DAB derivatives having new substituents at C2.

Scheme 8


Scheme 9


Alternatively, baccatin III may be used as a starting material and reacted as shown in Reaction Scheme 10.

5 After being protected at C7 and C13, baccatin III is reduced with LAH to produce 1,2,4,10 tetraol **24**. Tetraol **24** is converted to carbonate **25** using Cl_2CO and pyridine, and carbonate **25** is acylated at C10 with an acid chloride and pyridine to produce carbonate **26** (as shown) or with acetic anhydride and pyridine (not shown). Acetylation of carbonate

10 **26** under vigorous standard conditions provides carbonate **27** which is then reacted with alkyl lithiums to provide the baccatin III derivatives having new substituents at C2 and C10.


Scheme 10






9-desoxo-10-desacetoxy derivatives of baccatin III
 and 9-desoxo-10-desoxy derivatives of 10-DAB may be prepared
 by reacting baccatin III or 10-DAB (or their derivatives)
 5 with samarium diiodide and thereafter reducing the C9 keto
 substituent as otherwise described herein. Reaction between
 the tetracyclic taxane having a C10 leaving group and
 samarium diiodide may be carried out at 0°C in a solvent such
 as tetrahydrofuran. Advantageously, the samarium diiodide
 10 selectively abstracts the C10 leaving group; C13 side chains
 and other substituents on the tetracyclic nucleus remain
 undisturbed.

C7 dihydro and other C7 substituted taxanes can be
 prepared as set forth in Reaction Schemes 11 and 12.


REACTION SCHEME 11

REACTION SCHEME 12

As shown in Reaction Scheme 12, Baccatin III may be converted into 7-fluoro baccatin III by treatment with diethylaminosulfur trifluoride (DAST) at room temperature in THF solution. Other baccatin derivatives with a free C7 hydroxyl group behave similarly. Alternatively, 7-chloro baccatin III can be prepared by treatment of baccatin III with methane sulfonyl chloride and triethylamine in methylene chloride solution containing an excess of triethylamine hydrochloride.

A wide variety of tricyclic taxanes are naturally occurring, and through manipulations analogous to those described herein, an appropriate side chain can be attached to the C13 oxygen of these substances. Alternatively, as shown in Reaction Scheme 13, 7-O-triethylsilyl baccatin III

can be converted to a tricyclic taxane through the action of trimethyloxonium tetrafluoroborate in methylene chloride solution. The product diol then reacts with lead tetraacetate to provide the corresponding C4 ketone. This
5 ketone can be reduced to the alcohol with a hydride reducing agent such as sodium borohydride and subsequent acetylation produces the C4, C5 diacetate.

REACTION SCHEME 13

The following examples are provided to more fully illustrate the invention.

EXAMPLE 1

Taxol Using Zinc Alkoxide

5 To a solution of 7-triethylsilyl baccatin III (100 mg, 0.143 mmol) in 1 mL of THF at -45 °C was added dropwise 0.286 mL of a 0.5 M solution of potassium hexamethyldisilazide in toluene. After 15 min, 32 mg (0.143 mmol) of zinc chloride dimethoxyethane complex was added. After an
10 additional 1 h at -45 °C, the mixture was warmed to 0 °C and a solution of (+)-cis-1-benzoyl-3-triethylsilyloxy-4-phenylazetidin-2-one (82 mg, 0.215 mmol) in 1 mL of THF was added dropwise to the mixture. The solution was stirred at 0 °C for 3 h before 1 mL of a 10% solution of AcOH in THF was
15 added. The mixture was partitioned between saturated aqueous NaHCO₃ and 60/40 ethyl acetate/hexane. Evaporation of the organic layer gave a residue which was purified by flash chromatography followed by recrystallization to give 139 mg (90%) of (2'R,3'S)-2',7-(bis)triethylsilyl taxol.

20 EXAMPLE 2

Taxol Using Cadmium Alkoxide

 To a solution of 7-triethylsilyl baccatin III (100 mg, 0.143 mmol) in 1 mL of THF at -45 °C was added dropwise 0.286 mL of a 0.5M solution of potassium hexamethyldisilazide
25 in toluene. After 15 min, 26 mg (0.143 mmol) of anhydrous cadmium chloride was added. After an additional 1 h at -45 °C, the mixture is warmed to 0 °C and a solution of (+)-cis-1-benzoyl-3-triethylsilyloxy-4-phenylazetidin-2-one (82 mg, 0.215 mmol) in 1 mL of THF was added dropwise to the mixture.
30 The solution was stirred at 0 °C for 3 h before 1 mL of a 10%

solution of AcOH in THF is added. The mixture was partitioned between saturated aqueous NaHCO₃ and 60/40 ethyl acetate/ hexane. Evaporation of the organic layer gave a residue which was purified by flash chromatography followed by recrystallization to give 131 mg (85%) of (2'R,3'S)-2',7-(bis)triethylsilyl taxol.

EXAMPLE 3

Taxol Using Tetrabutylammonium Alkoxide

To a solution of 7-triethylsilyl baccatin III (100 mg, 0.143 mmol) in 1 mL of THF at -45 °C was added dropwise 0.286 mL of a 0.5 M solution of potassium hexamethyldisilazide in toluene. After 15 min, a solution of 16 mg (0.143 mmol) of anhydrous tetramethylammonium chloride in 0.5 mL of THF was added. After an additional 1 h at -45 °C, the mixture was warmed to 0 °C and a solution of (+)-cis-1-benzoyl-3-triethylsilyloxy-4-phenylazetidin-2-one (82 mg, 0.215 mmol) in 1 mL of THF was added dropwise to the mixture. The solution was stirred at 0 °C for 3 h before 1 mL of a 10% solution of AcOH in THF was added. The mixture was partitioned between saturated aqueous NaHCO₃ and 60/40 ethyl acetate/ hexane. Evaporation of the organic layer gave a residue which was purified by flash chromatography followed by recrystallization to give 134 mg (87%) of (2'R,3'S)-2',7-(bis)triethylsilyl taxol.